arXiv:1211.2837v1 [hep-ph] 12 Nov 2012

Left-Right Symmetry: from Majorana to Dirac

Miha Nemevšek,[1, 2] Goran Senjanović,[1] and Vladimir Tello[3, 4]
[1] ICTP, Trieste, Italy
[2] Jožef Stefan Institute, Ljubljana, Slovenia
[3] SISSA, Trieste, Italy
[4] Arnold Sommerfeld Center, LMU, München, Germany

(Dated: November 14, 2012)

Probing the origin of neutrino mass by disentangling the seesaw mechanism is one of the central issues of particle physics. We address it in the minimal left-right symmetric model and show how the knowledge of light and heavy neutrino masses and mixings suffices to determine their Dirac Yukawa couplings. This in turn allows one to make predictions for a number of low energy phenomena, such as neutrinoless double beta decay, electric dipole moments of charged leptons and neutrino transition moments.

I. Introduction. In the Standard Model (SM) all particles get their masses from the vacuum. This profound mechanism can be verified through the decays of the Higgs-Weinberg boson [1, 2], apparently found by CMS and ATLAS [3]. In particular, to every charged fermion of mass m_f corresponds a unique Yukawa coupling, which implies the following branching ratio

$$\Gamma\left(h \to f\overline{f}\right) \propto m_f^2. \tag{1}$$

What about neutrinos? Being neutral, they could be described by real Majorana spinors [4], as e.g., when one adds right-handed (RH) neutrinos to the SM [5]. However, even if one were able to measure both light and heavy neutrino masses and mixings, the Dirac couplings still could not be unambiguously determined [6].

The question is what happens in a more fundamental theory, such as the left-right (LR) symmetric model, introduced in order to understand the origin of parity violation [7]. Historically, this model led to neutrino masses long before the experiment and also to the seesaw mechanism [8, 9].

We show that once the mass matrix of heavy neutrinos is measured, the relation between heavy and light neutrinos can be made definite in the following manner. The KS [10] production process of heavy neutrinos allows one to measure their masses and flavour composition and determine their Majorana nature [11]. The theory then predicts the Dirac Yukawa couplings, shedding light on neutrinoless double beta decay and lepton dipole moments.

II. The Minimal LR Model. The minimal left-right symmetric model (LRSM) is based on the gauge group $SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$, augmented by a LR symmetry which implies equality of gauge couplings $g_L = g_R \equiv g$. Fermions come in LR symmetric doublet representations $Q_{L,R} = (u,d)_{L,R}$ and $L_{L,R} = (\nu, \ell)_{L,R}$ and the relevant charged gauge interactions are

$$\mathcal{L}_{gauge} = \frac{g}{\sqrt{2}}\left(\overline{\nu}_L V_L^\dagger \not{W}_L e_L + \overline{N}_R V_R^\dagger \not{W}_R e_R\right) + \text{h.c.}. \tag{2}$$

The Higgs sector consists [8] of a complex bi-doublet $\Phi(2,2,0)$ and two triplets $\Delta_L(3,1,2)$ and $\Delta_R(1,3,2)$ with quantum numbers referring to the LR gauge group.

In the seesaw picture the Majorana neutrino mass matrix is given by [12]

$$M_\nu = M_L - M_D^T \frac{1}{M_N} M_D, \tag{3}$$

where M_D is the neutrino Dirac mass matrix, while M_L and M_N are the symmetric Majorana mass matrices of left- and right-handed neutrinos, respectively.

The above formula connects the smallness of neutrino mass to the scale of parity restoration at high energies. It is crucial that there be new physical phenomena that allow to probe directly[1] the Majorana nature of RH neutrinos and determine their masses and mixings from experiment [10].

We opt for charge conjugation $\mathcal{C}$ as LR symmetry, with the fields transforming as $f_L \leftrightarrow (f_R)^c$, $\Phi \to \Phi^{\mathrm{T}}$ and $\Delta_L \leftrightarrow \Delta_R^*$ (the case of parity will be discussed elsewhere). The mass matrices then satisfy[2]

$$M_L = \frac{v_L}{v_R} M_N, \tag{4}$$

$$M_D = M_D^T, \tag{5}$$

where $v_{L,R}$ are the vevs of $\Delta^0_{L,R}$. In order to complete our phenomenological program, one needs to determine also v_R and v_L. The former would be automatically determined with the discovery of W_R; for the complex issues related to the latter we refer the reader to [14]. It is noteworthy that in the case of $\mathcal{C}$, the lower bound on the LR scale $M_{W_R} \gtrsim 2.5$ TeV [15, 16] is finally being probed at the LHC [17].

The determination of the RH neutrino mass matrix as a function of the Dirac Yukawa coupling (the latter imagined to be given by some external structure) was studied before in [18, 19]. We wish to show, on the contrary, that the LRSM is a complete theory of neutrino masses and mixings, in the sense that the measurements of the heavy sector at colliders can determine and inter-connect the low energy phenomena.

---

[1] In case RH neutrinos are too light to be probed at the LHC, one may still determine indirectly their masses and mixings as in the case when the lightest one is the warm dark matter [13].

[2] In principle, Φ could change sign under the LR symmetry. In case of $\mathcal{C}$, this implies anti-symmetric matrices with one vanishing and two other masses equal, clearly incompatible with data.

III. From Majorana to Dirac. The above seesaw formula seemingly obfuscates the connection between heavy and light neutrinos and common lore was that this connection cannot be unravelled [6]. However, since the Dirac mass matrix must be symmetric, it can be obtained directly from (3)

$$M_D = M_N \sqrt{\frac{v_L}{v_R} - \frac{1}{M_N} M_\nu}, \tag{6}$$

and thereby one can determine the mixing between light and heavy neutrinos. The square root of an n-dimensional matrix always has 2^n discrete solutions which can be found in [20] (ambiguities might arise in singular points of the parameter space). The essential fact is that Dirac masses are small, just as one would expect naively from the one generation case.

The above expression offers a unified picture of the low energy phenomena such as lepton flavour violation, lepton number violation through the neutrinoless double beta decay, electric dipole moments of charged leptons, neutrino transition moments, neutrino oscillations and neutrino cosmology. Some examples are discussed below, while the rest will be dealt with in a forthcoming publication.

M_N from LHC. The mass matrix of light neutrinos

$$M_\nu = V_L^* m_\nu V_L^\dagger \tag{7}$$

is being probed by low energy experiments, while the one of heavy neutrinos[3]

$$M_N = V_R m_N V_R^T \tag{8}$$

on the other hand, can be determined at high energy colliders through the KS reaction [10]. This amounts to producing W_R at the usual Drell-Yan resonance. Unlike in the case of W_L, where neutrinos act as missing energy, here the decays of heavy RH neutrinos lead to a lepton number violating final state of two same-sign leptons and two jets. Moreover, one can directly probe the Majorana nature of RH neutrinos through their equal branching ratios into charged leptons and anti-leptons [10]. Due to the absence of missing energy in the final state, one can fully reconstruct the heavy neutrino masses m_N and their mixings V_R by tagging the flavour of the final state leptons. For recent studies, see [15, 21].



FIG. 1. Electron EDM size in the LRSM with Eqs. (9), $v_L = 0$ and $m_{N_{1,2,3}} = 0.5, 2, 2.5$ TeV. The neutrino mixing angles are fixed at central values provided in [28] and the CP phases are scanned over.

While waiting for the LHC to provide this information, the reader may find it useful to have a simple working example

$$V_R = V_L^*. \tag{9}$$

Although in general (6) may require some computational tedium, for this example one gets

$$M_D = V_L^* m_N \sqrt{\frac{v_L}{v_R} - \frac{m_\nu}{m_N}} V_L^\dagger. \tag{10}$$

IV. Phenomenological implications. Low scale LRSM contains a host of experimentally accessible phenomena related to lepton number and flavour violation [22], both at high and low energies. In this section, we investigate only the low energy signals proportional to Dirac couplings and discuss their potential observation.

Electron EDM. One of the most sensitive probes of new physics beyond the SM is the T and CP-violating electric dipole moment (EDM) of charged leptons. The SM contribution arises at four loops [23] and is around eleven orders of magnitude below the current experimental limit $d_e < 10^{-27}$ e cm [24]. In the LRSM this process is significantly enhanced due to the mixing ξ_{LR} of left and right gauge bosons. The leading amplitude is present at one loop [25, 26]

$$d_e = \frac{e G_F}{4\sqrt{2}\pi^2} \mathrm{Im}\left[\xi_{LR} V_R F(t) V_R^\dagger M_D\right]_{ee}, \tag{11}$$

where

$$F(t) = \frac{t^2 - 11t + 4}{2(t-1)^2} + \frac{3t^2 \log t}{(t-1)^3}, \quad t = \frac{m_N^2}{M_{W_L}^2}. \tag{12}$$

There are strong limits on the ξ_{LR} but, in any case, it is automatically small due to the suppression of the heavy gauge boson mass. It is bounded by

$$\frac{\alpha}{4\pi} \frac{m_t m_b}{M_{W_R}^2} \lesssim |\xi_{LR}| \lesssim \frac{M_{W_L}^2}{M_{W_R}^2}, \tag{13}$$

---

[3] The mass matrix of charged leptons, being symmetric, can be taken diagonal without loss of generality.

with a lower bound resulting from radiative electroweak corrections [27].

Taking the example in (9), the size of the EDM is shown in Fig. 1 as a function of the lightest neutrino mass for two different neutrino hierarchies. These values can be probed by future experiments [29]. In the case when the LR mixing is close to its lower bound, one has to go beyond the one loop approximation [30], but in that case the experimental outlook seems bleak and we do not pursue it here.

In the context of LRSM, EDM is a manifestly CP odd process sensitive to Majorana phases, complementary to [31].

Neutrinoless double beta decay. The importance of this textbook example of lepton number violation was recognized in [32] soon after the seminal work of Majorana [4]. The LRSM offers new sources for this process [12] that has been studied extensively over the years [33]. In particular, an in-depth analysis [34] (see also [35]) was recently performed on the connection between neutrinoless double beta decay (and lepton flavour violation) at low energies and the KS process [10] at colliders.

Although the main source of this process in LRSM is due to the exchange of the heavy neutrinos, there is an additional contribution proportional to the Dirac mass matrix. We express it in the usual form of an effective mass term

$$m^{ee}_{\nu N} = \left(\xi_{LR} + \eta \frac{M^2_{W_L}}{M^2_{W_R}}\right) p \left(M_N^{-1} M_D\right)_{ee}, \tag{14}$$

where $p \simeq 100$ MeV [34] and $\eta \simeq 10^{-2}$ [36] are determined by nuclear physics considerations.

As a consequence of (6), the contribution in (14) is subleading[4] and the total decay rate is governed by the effective mass parameter

$$|m^{ee}_{\nu+N}|^2 = \left|V^2_{Lej} m_{\nu_j}\right|^2 + \left|p^2 \frac{M^4_W}{M^4_{W_R}} \frac{V^2_{Rej}}{m_{N_j}}\right|^2. \tag{15}$$

Since $|m^{ee}_{\nu+N}|$ and the size of the electron EDM both depend on the heavy neutrino mass, there is a correlation between the two processes, which is shown in Fig. 2. The values of m_N are chosen for illustration purpose only.

Neutrino transition moments. By defining

$$M = \xi^*_{LR} V_L^T m_\ell M_N^{-1} M_D V_L, \tag{16}$$

we get the following matrix of neutrino magnetic transition moments

$$\mu = \frac{ieG_F}{\sqrt{2}\pi^2} \mathrm{Im}\left[M + M^\dagger\right]. \tag{17}$$



FIG. 2. Electron EDM size and the effective mass for neutrinoless double beta computed with the same choice of parameters as in Fig. 1. The mass of W_R is fixed at 4 TeV and $p = 193$ MeV [34].

This result was already derived in [38] and can be reproduced using [26]. Here we neglect the contribution from light neutrino masses, which is roughly nine orders of magnitude smaller than the current experimental limit $\mu < 2 \times 10^{-10} \mu_B$ [39]. One should keep in mind that the Majorana transition moments in the SM (with non-zero neutrino mass) are negligibly small $\mu_{SM} \simeq 10^{-23} \mu_B$ [39].

It is easy to see that (17) gives roughly $\mu \simeq 10^{-19} \mu_B$ for generic values of M_D in (6), still a hopelessly small value. Therefore an observation of neutrino transition moments would deal a serious blow to the LRSM.

Alternative description. It is well known that in the SM augmented by RH neutrinos, the Dirac mass matrix is undetermined up to an arbitrary complex orthogonal matrix O [6, 40]. In the limit when the imaginary components of the mixing angles are large, the origin of neutrino mass is completely hidden from the processes that could probe it. The crucial point here is that M_D is forced to be symmetric, which now fixes O to have entries of natural order one. Moreover, this parametrization offers an alternative method of computing M_D which will be discussed elsewhere.

V. Conclusions and Outlook. In the SM the knowledge of charged fermion masses uniquely predicts Higgs decay branching ratios. As shown here, exactly the same happens in the LRSM for the masses of light and heavy neutrinos. The reason behind this is the LR symmetry itself which allows one to compute the Dirac Yukawa couplings in the context of the seesaw mechanism.

The main result of our paper is summarised in Eq. (6). Its phenomenological impact is exemplified on the phenomena of neutrinoless double beta decay, dipole moments of charged leptons and neutrino transition moments. This result was achieved at no expense of imposing additional ad-hoc symmetries but by the structure of the theory itself.

Acknowledgments. We thank A. Melfo for the inspirational version of the first abstract and F. Nesti for the early stages of collaboration and an important suggestion. We thank B. Bajc, A. Melfo and F. Nesti for a

---

[4] In addition, a possible contribution of heavy neutrinos without W_R [37] is suppressed according to (6).

careful reading of the manuscript. Also, we would like to thank K. Babu and Y. Zhang for useful discussions and comments. We acknowledge a continuing warm hospitality of BIAS. Special thanks are due to Ivo and Sonja Brzović for their kind hospitality, care and encouragement. V.T. is deeply grateful to Deniz for support.

---

[1] To the best of our knowledge, the existence of this boson was discussed during the 1964 only in P. W. Higgs, Phys. Rev. Lett. **13** (1964) 508.

[2] The precise properties (including the essential Yukawa couplings) of this particle were spelled out in S. Weinberg, Phys. Rev. Lett. **19** (1967) 1264.

[3] G. Aad *et al.* [ATLAS Collaboration], Phys. Lett. B **716** (2012) 1. [arXiv:1207.7214 [hep-ex]]; S. Chatrchyan *et al.* [CMS Collaboration], Phys. Lett. B **716** (2012) 30. [arXiv:1207.7235 [hep-ex]].

[4] E. Majorana, N. Cim. **14** (1937) 171.

[5] For a comprehensive review on neutrino mass, see: R. N. Mohapatra and A. Y. Smirnov, Ann. Rev. Nucl. Part. Sci. **56** (2006) 569 [hep-ph/0603118]; A. Strumia and F. Vissani, hep-ph/0606054.

[6] J. A. Casas and A. Ibarra, Nucl. Phys. B **618** (2001) 171 [hep-ph/0103065].

[7] J.C. Pati, A. Salam, Phys. Rev. D **10** (1974) 275; R. N. Mohapatra, J.C. Pati, Phys. Rev. D **11** (1975) 2558; G. Senjanović, R. N. Mohapatra, Phys. Rev. D **12** (1975) 1502; G. Senjanović, Nucl. Phys. B **153** (1979) 334.

[8] P. Minkowski, Phys. Lett. B **67** (1977) 421; R. N. Mohapatra, G. Senjanović, Phys. Rev. Lett. **44** (1980) 912.

[9] T. Yanagida, *Workshop on unified theories and baryon number in the universe*, ed. A. Sawada, A. Sugamoto (KEK, Tsukuba, 1979); S. Glashow, *Quarks and leptons, Cargèse 1979*, ed. M. Lévy (Plenum, NY, 1980); M. Gell-Mann *et al.*, *Supergravity Stony Brook workshop, New York, 1979*, ed. P. Van Niewenhuizen, D. Freeman (North Holland, Amsterdam, 1980).

[10] W.-Y. Keung, G. Senjanović, Phys. Rev. Lett. **50** (1983) 1427.

[11] For recent reviews, see: G. Senjanović, Int. J. Mod. Phys. A **26** (2011) 1469 [arXiv:1012.4104 [hep-ph]]; G. Senjanović, Riv. Nuovo Cim. **034**, 1 (2011).

[12] R. N. Mohapatra, G. Senjanović, Phys. Rev. **D23** (1981) 165.

[13] M. Nemevšek, G. Senjanović and Y. Zhang, JCAP **1207** (2012) 006 [arXiv:1205.0844 [hep-ph]]; F. Bezrukov, H. Hettmansperger and M. Lindner, Phys. Rev. D **81**, 085032 (2010) [arXiv:0912.4415 [hep-ph]].

[14] An non exhaustive list of references: P. Fileviez Perez, T. Han, G. -y. Huang, T. Li and K. Wang, Phys. Rev. D **78** (2008) 015018 [arXiv:0805.3536 [hep-ph]]; A. Melfo, M. Nemevšek, F. Nesti, G. Senjanović and Y. Zhang, Phys. Rev. D **85** (2012) 055018 [arXiv:1108.4416 [hep-ph]]; S. Godfrey and K. Moats, Phys. Rev. D **81** (2010) 075026 [arXiv:1003.3033 [hep-ph]]; C. -W. Chiang, T. Nomura and K. Tsumura, Phys. Rev. D **85** (2012) 095023 [arXiv:1202.2014 [hep-ph]].

[15] A. Maiezza, M. Nemevšek, F. Nesti and G. Senjanović, Phys. Rev. D **82** (2010) 055022; [arXiv:1005.5160 [hep-ph]].

[16] Y. Zhang, H. An, X. Ji and R. N. Mohapatra, Nucl. Phys. B **802** (2008) 247 [arXiv:0712.4218 [hep-ph]]; M. Blanke, A. J. Buras, K. Gemmler and T. Heidsieck, JHEP **1203** (2012) 024 [arXiv:1111.5014 [hep-ph]]; S. Bertolini, J. O. Eeg, A. Maiezza and F. Nesti, arXiv:1206.0668 [hep-ph].

[17] S. Chatrchyan *et al.* [CMS Collaboration], arXiv:1210.2402 [hep-ex]; G. Aad *et al.* [ATLAS Collaboration], Eur. Phys. J. C **72** (2012) 2056 [arXiv:1203.5420 [hep-ex]].

[18] D. Falcone, Phys. Rev. D **68** (2003) 033002 [hep-ph/0305229].

[19] E. K. Akhmedov and M. Frigerio, Phys. Rev. Lett. **96** (2006) 061802 [hep-ph/0509299]; P. Hosteins, S. Lavignac and C. A. Savoy, Nucl. Phys. B **755** (2006) 137 [hep-ph/0606078]; E. K. Akhmedov and M. Frigerio, JHEP **0701** (2007) 043 [hep-ph/0609046].

[20] Gantmacher F. R., The theory of matrices, Vol. 1. Chelsea Pub. Co., New York, (1960).

[21] M. Nemevšek, F. Nesti, G. Senjanović and Y. Zhang, Phys. Rev. D **83** (2011) 115014 [arXiv:1103.1627 [hep-ph]]; C. -Y. Chen and P. S. B. Dev, Phys. Rev. D **85** (2012) 093018 [arXiv:1112.6419 [hep-ph]]; S. P. Das, F. F. Deppisch, O. Kittel and J. W. F. Valle, Phys. Rev. D **86** (2012) 055006 [arXiv:1206.0256 [hep-ph]]; J. A. Aguilar-Saavedra and F. R. Joaquim, arXiv:1207.4193 [hep-ph].

[22] V. Cirigliano, A. Kurylov, M. J. Ramsey-Musolf and P. Vogel, Phys. Rev. Lett. **93** (2004) 231802 [hep-ph/0406199].

[23] M. Pospelov and A. Ritz, Annals Phys. **318** (2005) 119 [hep-ph/0504231].

[24] J. J. Hudson, D. M. Kara, I. J. Smallman, B. E. Sauer, M. R. Tarbutt and E. A. Hinds, Nature **473** (2011) 493.

[25] An original calculation in the context of LRSM can be found in J. F. Nieves, D. Chang and P. B. Pal, Phys. Rev. D **33** (1986) 3324.

[26] For a complete general study of this and similar one loop transitions, see: L. Lavoura, Eur. Phys. J. C **29** (2003) 191 [hep-ph/0302221].

[27] G. C. Branco and G. Senjanović, Phys. Rev. D **18** (1978) 1621.

[28] M. C. Gonzalez-Garcia, M. Maltoni, J. Salvado and T. Schwetz, arXiv:1209.3023 [hep-ph].

[29] http://electronedm.org

[30] D. Atwood, C. P. Burgess, C. Hamzaoui, B. Irwin and J. A. Robinson, Phys. Lett. B **256** (1991) 471.

[31] A. de Gouvea, B. Kayser and R. N. Mohapatra, Phys. Rev. D **67** (2003) 053004 [hep-ph/0211394].

[32] G. Racah, N. Cim. **14** (1937) 322; W. H. Furry, Phys. Rev. **54** (1938) 56; W.H. Furry, Phys. Rev. **56** (1939) 1184.

[33] For a recent review, see: W. Rodejohann, Int. J. Mod. Phys. E **20**, 1833 (2011) [arXiv:1106.1334 [hep-ph]].

[34] V. Tello, M. Nemevšek, F. Nesti, G. Senjanović and F. Vissani, Phys. Rev. Lett. **106** (2011) 151801 [arXiv:1011.3522 [hep-ph]]; M. Nemevšek, F. Nesti, G. Senjanović and V. Tello, arXiv:1112.3061 [hep-ph].

[35] J. Chakrabortty, H. Z. Devi, S. Goswami and S. Patra, JHEP **1208** (2012) 008 [arXiv:1204.2527 [hep-ph]].

[36] M. Doi, T. Kotani and E. Takasugi, Prog. Theor. Phys. Suppl. **83** (1985) 1; K. Muto, E. Bender and H. V. Klapdor, Z. Phys. A **334** (1989) 187; M. Hirsch, H. V. Klapdor-Kleingrothaus and O. Panella, Phys. Lett. B **374** (1996) 7 [hep-ph/9602306].

[37] A. Atre, T. Han, S. Pascoli and B. Zhang, JHEP **0905** (2009) 030 [arXiv:0901.3589 [hep-ph]]; M. Mitra, G. Senjanović and F. Vissani, Nucl. Phys. B **856** (2012) 26 [arXiv:1108.0004 [hep-ph]]; M. Mitra, G. Senjanović and F. Vissani, arXiv:1205.3867 [hep-ph]; J. Lopez-Pavon, S. Pascoli and C. -f. Wong, arXiv:1209.5342 [hep-ph].

[38] M. Czakon, J. Gluza and M. Zralek, Phys. Rev. D **59** (1999) 013010.

[39] C. Broggini, C. Giunti and A. Studenikin, arXiv:1207.3980 [hep-ph].

[40] E. K. Akhmedov and W. Rodejohann, JHEP **0806** (2008) 106 [arXiv:0803.2417 [hep-ph]].